



Square-shaped Alopecia After Embolization of Intracranial Aneurysm: a Case Report and Review

Jorge Román-Sainz, Nicolás Silvestre-Torner, Fernando Gruber-Velasco,
Belen Romero-Jiménez, Alejandro Lobato-Berezo, Adrián Imbernón-Moya

Author	Age	Gender	Radiation (Gy)	Biopsy	Latency (weeks)	Resolution (months)	Treatment
S. Podlipnik	38	M	ED: >4	No	4	2	No
W. Huda	34	F	6,6	No	NS	3	NS
G. M. Garnacho	39	M	ED: 3	No	2	5	No
A. César	26	F	5	Yes	1	3	No
M. Freysz (18 P)	AR: 19-58	6 M, 12F	2,5-12,3	No	2-4	2-3	No
M. D'incan (6 P)	AR: 27-47	4 M, 2 F	NS	No	2	3-4	No
M. Krasovec	32	M	NS	No	2	3	No
N. Marti	29	F	ED >3	No	2	4	No
WS. Lee (5 P)	AR: 40-57	1 M, 4 F	ED >2	Yes	2	5-6	Cryotherapy and deoxymethasone
V. López	44	F	ED >3	No	3	4	No
CS. Wen	33	M	ED >3	No	2	2	NS
R. Nannapaneni (2 P)	AR: 31-45	2 F	3,83-4,07	No	4	6	NS
JE Seol (3 P)	AR: 33-51	2 M, 1 F	NS	Yes	2	3-4	Minoxidil and topical corticosteroids and triamcinolone.
S. Verma	47	F	ED >3	No	2	3	No
M. Vaccaro ²	NS	F	3,92	No	4	2	No
A. Vazquez	13	M	NS	No	6	6	No
S Cho ³ (10 P)	AR: 35-70	2 M, 8 F	NS	Yes	1-8	3-4	No
H Lee (2 P)	AR: 49-55	2 M	4,2-5,7	No	2	4-5	Topical minoxidil
C Bernárdez	32	F	NS	Yes	4	2	No

P: Patients AR: Age Range M: Male F: Female ED: Estimated Dose NS: Not Specified