

Amelanotic Melanoma Presenting As a Scaly Plaque

Catalina Aitken¹, Natalia Gómez², Cristian Navarrete-Dechent¹, Andrea Antúnez-Lay^{1,3}

¹ Department of Dermatology, Pontificia Universidad Católica de Chile, Santiago, Chile

² Department of Pathology, Hospital Dr. Sotero del Río, Santiago, Chile

³ Department of Dermatology, Hospital Dr. Sotero del Río, Santiago, Chile

Citation: Aitken C, Gómez N, Navarrete-Dechent C, Antúnez-Lay A. Amelanotic melanoma presenting as a scaly plaque. *Dermatol Pract Concept.* 2026;16(2):6348. DOI: <https://doi.org/10.5826/dpc.1602a6348>

Accepted: August 28, 2025; **Published:** April 2026

Copyright: ©2026 Aitken et al. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (BY-NC-4.0), <https://creativecommons.org/licenses/by-nc/4.0/>, which permits unrestricted noncommercial use, distribution, and reproduction in any medium, provided the original authors and source are credited.

Funding: None.

Competing Interests: None.

Authorship: All authors have contributed significantly to this publication.

Corresponding Author: Catalina Aitken Bravo, Resident of Dermatology, Department of Dermatology, Escuela de Medicina, Pontificia Universidad Católica de Chile, 4686 Vicuña Mackenna Avenue, Santiago, Chile. E-mail: caitkenb@gmail.com

Case Presentation

An 86-year-old female with a history of multiple non-melanoma skin cancers and extensive sun damage was evaluated for an eczematous lesion on the left arm. Dermatological examination revealed a 40 x 58 mm pink scaly plaque with structureless tan areas, shiny white lines, and a solitary blue-grey globule under dermoscopy. An excisional biopsy was performed that confirmed a 0.5-mm depth invasive amelanotic melanoma (AM).

Teaching Point

The diagnosis of AM remains a challenge. It usually presents as papulonodular or ulcerated lesions (58%) [1]. Other rare forms can present as erythematous macules or patches or as eczema-like plaque as in our case [2]. They lack 'ABCD'

criteria, and due to the lack of pigmentation, there is a diagnosis delay [1,2]. When it comes to solitary eczematous lesions resistant to topical treatment, the differential diagnosis includes basal cell carcinoma (BCC), Bowen's disease, extramammary Paget's disease, lichen sclerosus et atrophicus and, less frequently, mycosis fungoides. Although a scaly plaque being a rare presentation, AM should also be included in this differential, especially in patients with extensive sun damage and a history of skin malignancy. Dermoscopy can provide important clues, including the presence of dotted, serpentine, and polymorphous vessels, milky-red areas, shiny white structures, and light brown structureless areas (in hypomelanotic melanomas) [2]. In this case, the presence of polymorphous vessels and structureless tan areas should raise the possibility of AM; however, the scale and 'orangy' areas were more suggestive of eczema, and the presence of a 'bluish globule' misled us towards BCC.




Figure 1. Amelanotic melanoma. A: Clinical photo showing a pink, ill-defined, scaly plaque. B: Dermoscopic photo showing tan structureless areas (black arrows) and polymorphous vessels (red arrows) (polarized light dermoscopy, original magnification 10x). C: Dermoscopic photo showing a “blue-grey globule” (white arrow) and polymorphous vessels (red arrow) (polarized light dermoscopy, original magnification 10x). D: Photomicrograph showing an atypical proliferation of melanocytes in the junction with pagetoid ascent (H&E, 20x).

References

1. Gong HZ, Zheng HY, Li J. Amelanotic melanoma. *Melanoma Res.* 2019;29(3):221-230. DOI:10.1097/CMR.0000000000000571
2. Jaimes N, Braun RP, Thomas L, Marghoob AA. Clinical and dermoscopic characteristics of amelanotic melanomas that are not of the nodular subtype. *J Eur Acad Dermatol Venereol.* 2012;26(5):591-596. DOI:10.1111/j.1468-3083.2011.04122.x