

Supplementary File

Table S1. Sample categorization according to NLR and age subgroups.

		ADULT 18-65 yo	OLDER ADULTS 65-75 yo	AGED ≥ 75 yo
Robust NLR≤2.44	M	92 (14.58%)	30 (4.75%)	137 (21.71%)
	F	47 (7.45%)	72 (11.41%)	41 (6.5%)
	Tot	139 (22.03%)	102 (16.16%)	178 (28.2%)
Frail 2.44-3.53 NLR	M	14 (2.22%)	24 (3.80%)	55 (8.72%)
	F	22 (3.49%)	21 (3.33%)	9 (1.43%)
	Tot	36 (5.71%)	45 (7.13%)	64 (10.14%)
Extremely frail NLR≥3.55	M	17 (2.69%)	10 (1.58%)	25 (3.96%)
	F	5 (0.79%)	1 (0.16%)	9 (1.43%)
	Tot	22 (3.49%)	11 (1.74%)	34 (5.39%)

Table S2. ESR levels according to age group. Data are indicated as Mean±SD and percentages.

ESR	ADULT 18-65 yo	OLDER ADULTS 65-75 yo	AGED ≥ 75 yo	Tot (n=117)
M	10.11 ± 9.73 (n=35; 29.91%)	14.22 ± 7.14 (n=9; 7.69%)	11.87 ± 7.27 (n=15; 12.82%)	11.18 ± 8.8 (n=59; 50.4%)
F	14.89 ± 9.39 (n=18; 15.38%)	12.95 ± 8.1 (n=20; 17.09%)	25.55 ± 21.26 (n=20; 17.09%)	17.9 ± 15.19 (n=58; 49.57%)
Tot	11.73 ± 9.79 (n=53; 45.3%)	13.34 ± 7.71 (n=29; 24.8%)	19.69 ± 17.93 (n=35; 29.91%)	14.51 ± 12.79 (n=117; 100%)

Table S3. CRP levels according to age group. Data are indicated as Mean±SD and percentages.

CRP	ADULT 18-65 yo	OLDER ADULTS 65-75 yo	AGED ≥ 75 yo	Tot (n=135)
M	2.95 ± 4.08 (n=32; 23.7%)	7.07 ± 12.23 (n= 21; 15.56%)	11.62 ± 29.54 (n=23; 17.04%)	6.71 ± 17.78 (n=76; 56.3%)
F	2.19 ± 2.1 (n=15; 11.1%)	2.73 ± 2.33 (n=27; 20%)	5.56 ± 10.18 (n=17; 12.6%)	3.41 ± 5.83 (n=59; 43.7%)
Tot	2.71 ± 3.57 (n=47; 34.82%)	4.63 ± 8.45 (n=48; 35.56%)	9.05 ± 23.32 (n=40; 29.63%)	5.27 ± 13.94 (n=135; 100%)